


UNCLASSIFIED

N00014-86-K-0289

NL

F/G 5/8

NL



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A184 058

DTIC FILE COPY

12

STUDIES OF THE PROCESSING OF SINGLE WORDS USING
POSITRON TOMOGRAPHIC MEASURES OF CEREBRAL BLOOD FLOW CHANGE

Steven E. Petersen, Peter T. Fox, Michael I. Posner, Marcus Raichle

McDonnell Center for Studies of Higher Brain Function
Washington University, St. Louis

ONR 87-7

DTIC
ELECTE
S AUG 25 1987 D
D

Research sponsored by: Personnel and Training Research Program,
Psychological Sciences Division
Office of Naval Research
Under Contract Number: N0014-86-K-0289
Contract Authority Number: NR-442a554

Reproduction in whole or part is permitted for any
purpose of the United States Government.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

87 8 21 004

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release: distribution unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report #87-7			5. MONITORING ORGANIZATION REPORT NUMBER(S) Personnel & Training Research Programs Office of Naval Research (CODE 1142PT)		
6a. NAME OF PERFORMING ORGANIZATION Washington University		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) 660 S. Euclid, Box 8111 Department of Neurology St. Louis, MO 63110			7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-KO289		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 61153N	PROJECT NO. RR04206	TASK NO. RR04206-0A
					WORK UNIT ACCESSION NO. NR442a554
11. TITLE (Include Security Classification) Studies of the Processing of Single Words Using Positron Emission Tomographic Measurements of Cerebral Blood Flow Change					
12. PERSONAL AUTHOR(S) Petersen, Steven E.					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 01MAY87 TO 01MAY88		14. DATE OF REPORT (Year, Month, Day) August 3, 1988	
15. PAGE COUNT 51					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
05	10				
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS					
21. ABSTRACT SECURITY CLASSIFICATION Unclassified					
22a. NAME OF RESPONSIBLE INDIVIDUAL Michael I. Posner			22b. TELEPHONE (Include Area Code) (314) 362-3317		22c. OFFICE SYMBOL ONR 1142 PT

STUDIES OF THE PROCESSING OF SINGLE WORDS USING AVERAGED POSITRON EMISSION TOMOGRAPHIC MEASUREMENTS OF CEREBRAL BLOOD FLOW CHANGE

Steven E. Petersen, Peter T. Fox, Michael I. Posner, Marcus E. Raichle

INTRODUCTION

Language is an essential characteristic of the human species, and as such has been a focal point for study in disciplines ranging from philosophy to neurology. The totality of language is amazingly complex and includes the study of syntax, semantics and pragmatics. Cognitive and neurological investigations of language often narrow the focus of study to the processing of individual words (lexical items). Lexical processing involves a network of several levels of internal coding that can be isolated by experiment. (LaBerge & Samuels, 1974; Rumelhart & McClelland, 1982, 1986). Examples of some of these separate codes include a visual image of the form of a spoken word (visual code), pronunciation of the word (phonological code) or the association of related words (semantic codes). Studies of the time course of activation of these internal codes of words and the roles they play in performance has been a central topic in the cognitive psychology of reading and listening. (Carr & Pollatsek, 1985; Posner, 1978 and Rumelhart & McClelland, 1986 for examples). Behavioral neurologists have been concerned as well with issues in lexical processing, but the focus has been in correlating the internal stages of processing with different brain regions. (Geschwind, 1979).

Recent advances in activation techniques and data analysis strategies using positron emission tomographic (PET) measurements of blood flow (BF) change have made it possible to address concerns relevant to both cognitive science and behavioral neurology. These techniques have been used to make precise anatomical-functional correlations in the primary and secondary sensory and motor areas of the human brain (e.g., the mapping of the visual field onto primary visual cortex, Fox, et al., 1987) by comparing several test and control conditions within a single session. In this paper, we take advantage of these techniques to study the neural mechanisms underlying the processing and production of single words in normal subjects. Our results indicate that the processing of single lexical items during simple sensory, word repetition, and word generation tasks activates a small number of discrete areas. These findings suggest a framework for the processing of single words that is quite different from the standard clinical neurological account based on cortical lesions (Geschwind, 1965; 1979), but is consistent with recent results from cognitive science (Coltheart, 1985; Carr & Pollatsek, 1985).

METHODS

PET techniques common to all studies

Tracer strategy Changes in neural activity induce changes in local BF (Fox & Raichle, 1986). We used O15-labeled water injected as an intravenous

bolus to measure BF in normal subjects. The short half life of ^{15}O (120 s) allowed a short interscan interval of 10 minutes (5 half lives) and acquisition of 8-10 40 sec scans per subject. This in turn allowed comparisons of several different task conditions within single subjects, who remain in the same position with respect to the scanner throughout the session.

Image subtraction Side-by-side visual inspection of the blood flow measurements in two conditions gives some information about the presence of focal blood flow changes (Fig 1A and B, page 10), but to attain quantitative information about the location and magnitude of differences between the scans, subtraction of the activity in one image from another (activated - control condition) were performed. When a direct pixel-by-pixel subtraction is made, the resultant image was of foci of change between the two primary images (Fig 1C). The subtractions were made after linear normalization that negated the effects of global BF fluctuation. The global brain blood flow was calculated using a standardized mean regional method (Perlmutter, et al, 1985). Each scan was normalized by multiplying every pixel of each scan by a correction factor, calculated as the mean normal global BF of this laboratory [$50 \text{ ml}/(100 \text{ gm} \cdot \text{min})$] divided by the scan global BF, before computing subtraction images.

Anatomical standardization Because the precise alignment and position of the seven PET slices vary from subject to subject, foci of change in subtraction images are in coordinates unique to the individual being scanned. To anatomically identify areas of change, each individual's scan information must be converted to a standardized reference space. To achieve this standardization, the seven slice information on each individual is first subjected to an interslice interpolation to fill an array of uniform pixel size (about 2 mm). These arrays were then standardized to a single anatomical coordinate space using information from a lateral skull X-ray taken while the subject was in position, and slice length and width information from the scans themselves. This anatomical standardization made it possible for the investigator to "look up" the anatomical location of the areas of change in a standard atlas (Fox, Perlmutter & Raichle, 1985).

Image averaging The anatomical standardization of subtraction scans to an individual anatomical standard gives a second advantage: subtraction images during the same condition from different individuals can be summed and averaged. The rationale behind image averaging is that areas of activation that are consistent across individuals will sum and those that are not consistent will be random and cancel one another out. This simple signal averaging technique greatly enhances the detection and localization of the subtle activations that are produced during cognitive manipulations (Mintun, Fox, and Raichle).

Automated response detection The subtraction images from single individuals and averaged subtraction images across individuals show many foci of change. A computer search routine was designed to find all local peaks of activity change within an image, and record their location and magnitude. This routine was used for objective localization of areas of change in activated pair subtractions (Fox, et al, 1986).

Statistical Analysis As the local-maximum search routine allowed all independent local changes to be identified within any subtraction image, significance testing was based upon analysis of this entire population. During the averaging process spatially nonrandom changes (i.e., stimulus-related responses) gained in magnitude relative to random changes (i.e., noise), lying progressively further (more standard deviations) from the mean. That is, consistent responses became "outliers" in the population. This prompted the application of tests for outlier detection as a means of identifying significant responses (Snedecor and Corcoran, 1980). Our statistical analysis was two-tiered, first using omnibus tests (g^1 and g^2 statistics) to determine whether a data set (a population of regional differences) contained significant outlier responses, then applying post-hoc analysis (Z-score) to determine the significance levels of specific responses within the population (see tables I-VII). All responses with Z scores giving a P-value less than .03 are reported.

Stimulus presentation Visual stimuli were presented on a color monitor suspended about 12 inches in front of the subject perpendicular to the line of sight. Auditory stimulation was generated through small speakers placed in the ear of the subject. The speakers were driven by a digital tape recorder on which appropriate stimuli have been recorded. All stimuli were single common nouns (e.g., cake, radio, etc.), and were presented at 1 Hz. The visual stimuli were presented for 150 ms with an 850 ms interword interval. The auditory stimuli varied in presentation time with the length of the word.

Subjects The subjects were medical students, medical residents, psychology graduate students and represented a high normal group for intelligence and reading skills. All were native English speakers and were right-handed as assessed with the Edinburgh handedness inventory.

Paradigm design The paradigms were designed to address two interrelated issues. 1) To utilize our battery of data analysis techniques, it is important to design runs in activated-control subtraction pairs. In these experiments, a stepwise design was used so that the stimulated member of one pair became the control for the next pair. 2) To make the images of change interpretable, an attempt must be made to hold all but a small number of task components in the two runs constant so that the changes in blood flow may be attributed to a small number of specific functions.

The lexical processing experiments presented here represent a first attempt to address some basic issues in the processing and production of single words (11). There were four basic tasks that the subjects were asked to perform; each task was performed twice, once with visual input and once with auditory input.

The core tasks were: 1) to fixate on a small crosshair with no other stimulation (fixation point); 2) passive nouns presented visually or auditorily through the speakers while the person fixates the crosshair (passive words); 3) repeating aloud the visually or auditorily presented nouns (repeat words); 4) generating aloud an appropriate verb or use for the presented noun, for example, to say the word "eat" if the presented word was "cake" (generate uses).

The core tasks then were organized into three stepwise subtractions. In the first level subtraction, activity during the fixation point condition was subtracted from the passive words condition (sensory subtraction). The second level was the subtraction of passive words condition from the repeat words condition (motor subtraction). The third level subtraction was of the repeat words condition from the generate uses condition (cognitive subtraction). Some other subtractions, spanning more than one level were performed to assess questions about our subtractive assumptions.

We recognize the difficulty of designing two tasks that are the same except for the addition of a single new operation (Sternberg, 1969). Indeed, the activations in our tasks are often due to more than one type of operation. It is also possible that subjects alter their basic strategy when confronted with a new task. However, by appropriate subtractions one can check up on whether strategies for more complex tasks still involve activation of those areas comprising simpler tasks. Some secondary experiments were also performed to aid in the interpretations of the results, and will be explained where appropriate.

RESULTS

Sensory Subtraction

A comparison of the resting task and the passive nouns task (sensory subtraction) showed modality specific activation in those areas that are related to the basic sensory information contained in the words, namely primary visual and auditory cortices. In addition, higher order, but modality specific areas were identified that appear to be specific to higher level, possibly lexical processing. There are no regions that are activated by the passive presentation of both auditory and visual words.

Visual (Passive visual words - fixation point) (PVW)

As would be expected, the most responsive foci of activation in the visual sensory subtraction are found in the occipital lobe. There is a bilateral response in buried calcarine striate cortex (primary visual cortex, foci 1, 2; Table 1). There are also extrastriate regions of response (Fig. 3): one region is represented bilaterally (Foci 3,4) with about 30% greater activation on the left, and a region still further anterior and inferior in right occipital cortex (Focus 5). This is also a left lateralized area of activation in the basal ganglia, probably in the putamen (Focus 6).

The occipital pole has long been associated with visual function so that occipital activation here is hardly surprising. However, in a recent PET activation studies, simple visual checkerboard (Fox, et al, 1986) or random dot stimuli (unpublished observations) did not produce consistent responses in lateral occipital cortex anterior to striate cortex, areas that are clearly activated during the presentation of visual words. The lateral occipital activation may well represent activation reflecting lexical or letter level processing (see discussion).

Auditory (Passive auditory words - fixation point) (PAW)

The presentation of auditory words activates several regions in and near the posterior temporal lobe. From animal and human studies, primary auditory cortex is presumed to be located in the posterior third of the temporal sylvian cortex (Geschwind & Levitsky, 1968; Imig, et al, 1977). We find a clear bilateral activation in this region (Fig. 4, Foci 7,8; Table II). There is also a second focus of activation, in the left hemisphere, anterior and inferior to the primary activation (Fig. 3, Focus 9, Table II), and this focus may represent one of the secondary auditory areas that have been described extensively in other species, for example, by Imig et al (1977). There is an active focus in inferior anterior cingulate cortex (Fig. 4, Focus 12, Table II). Finally, activation can also be found in the supramarginal gyrus (SMG, Fig. 4, Focus 10) in the left hemisphere. The SMG is the only activation focus in any of our tasks that is in the Wernicke's-angular gyrus region that is commonly associated with language processing. Since neither the inferior temporal (Focus 9) or supramarginal gyrus has been shown to be active for simple auditory stimuli, including tones, clicks, or rapidly-presented synthetic syllables (Lauter, et al, 1984, and unpublished observations), these foci could be considered as candidates for regions performing word-level processing.

Motor Subtraction

When the repeat task is compared to the passive task (motor subtraction), areas related to overt reading, the generation of an articulatory code, motor programming, and the generation of the movements themselves are activated. Since we are focusing on the output, which is similar across the two input modalities, and subtracting the different sensory specific responses, we would expect that the structures that would be activated would be similar for the auditory and visual presentations and our results confirm this expectation.

Visual (Repeat visual words - passive visual words) (RVW)

One area of activation in the repeat conditions, Rolandic cortex (Fig. 5, Foci 13, 14; Table III), represents primary motor cortical activation that would be expected to be present in the repeat condition. Rolandic cortex has been associated with somatosensory and motor function, primary somatosensory cortex located posteriorly and primary motor cortex located anteriorly in this region. Both of these representations are laid out somatotopically so that adjacent parts of the body are represented in adjacent parts of the cortex. Recent studies using vibrotactile stimulation have identified the locations in our stereotactic space of toes, hand and lips somatosensory representation (Fox, Burton & Raichle, 1987). The area identified as Rolandic mouth cortex is just anterior and superior to the somatosensory lips representation.

A second area activated in the repeat condition is superior, anterior cerebellum (Foci 19,20; Table III). The cerebellum has also been associated with aspects of motor coordination, and cerebellar activation during the speech output is expected. Other studies, where subjects performed hand and eye movements, have shown activation in a similar location (Fox, Thach & Raichle, 1987).

Speech production also bilaterally activated a region of the inferior frontal operculum anterior to the Rolandic strip (Fig. 6, Foci 15,16). The left site of activation is in a location often defined as Broca's area. Damage to Broca's area in the left hemisphere is often thought to be associated with speech production deficits and agrammatism. In our paradigms, however, the activation is bilateral, and while this area could be related to some specifically linguistic functions. Simple tongue movement (Fig. 7) and hand movement (Fox, Pardo, Petersen and Raichle, 1987) both cause activation in a similar location. Given these observations, it seems most plausible at this time to associate this region with some aspect of motor programming, or praxis, than with specifically linguistic or speech function.

A region of medial frontal cortex, in a location often identified as the supplementary motor area (SMA), is activated in the repeat visually presented words task (Fig. 8, Focus 18, Table III). A covert task which required a subject to monitor and count pairs of associated targets with no overt output during the task also activated this region (Petersen, et al, 1986, Soc. Neurosci.).

Finally, there is an activation focus (17, Table III) which lies anterior and inferior to the Rolandic mouth representation, that is left lateralized, and a focus of activation in the colliculus (Focus 21; Table III).

Auditory (Repeat auditory words - passive auditory words) (RAW)

The most startling aspect of the results from the RAW subtraction is the degree of similarity between the results from auditory and visual presentation. With only a small number of exceptions, foci activated when repeating auditorily presented words were also activated when repeating visually presented words, when the passive sensory activation is subtracted away.

The area in RAW overlapping RVW includes bilateral rolandic mouth representation (Foci 22,23; Table IV), left buried "Broca" cortex (Focus 24, Table V), right lateral "Broca's" cortex (Focus 25; Table IV), SMA (Focus 27, Table IV), and left premotor cortex (Focus 66, Table IV).

For most of these foci, the anatomical location between input modalities is within .5 cm. In the case of the left premotor area, however, the slightly larger difference between locations may well represent a modality specific specialization of adjacent zones having similar functions.

There are a small set of activation foci not shared across modalities including the superior, anterior cerebellum and colliculus (Foci 19,20, 21; Table III) for RPW.

Cognitive Subtraction

Our final comparison is the most subtle, and should represent activation related to purely cognitive processes, since the sensory input is identical in the stimulated and control condition, and the motor output is very similar. Again, since similar operations are called for in the performance

of the task for auditory or visual presentations, there were similar areas activated in the input conditions.

Visual (Generate visual words - Repeat visual words) (GVW)

Inferior and anterior to the SMA is a region activated in the generate uses tasks (Fig. 9, Focus 31, Table V). This area is located in the anterior cingulate cortex (ACC). Inferior and anterior to this midline activation is a second focus of anterior cingulate activation (Focus 32, Table V).

Left lateralized anterior inferior cortex in and around areas 45, 46, and 47 shows several active foci only during the generation task (Fig. 10, Foci 28, 29, 30; Table V). Our choice of a verb (use) associate to a noun confounds grammatical and semantic transformations. The lesion literature shows effects that appear to be related to verb association, and agrammatism associated with lesions in these regions (McCarthy & Warrington, 1985). Subjects with Broca's aphasia have deficits on semantic priming tasks (Milberg & Blumstein, 1981; Milberg, Blumstein & Dvoretzky, 1987). So, in a converging experiment, a semantic monitoring task (monitoring strings of nouns for members of a semantic category), an inferior frontal region near focus 35 was activated (see Fig. 11). Since one task involves monitoring for semantic association of nouns in a string of presented nouns, and the other involves the generation of a semantically-associated verb, it seems likely that some common computation involved in semantic association takes place in the anterior inferior frontal lobe. It remains to be seen whether syntactic operations, will also involve these foci.

There is also a striking focus of activation in right inferior lateral cerebellum (Fig. 12, Focus 36, Table V). The lateralization to the right cerebellar hemisphere is consistent with observations that a cerebellar hemisphere is anatomically and functionally related to the contralateral (in this case left language dominant) cerebral hemisphere. It seems unlikely that this represents motor activation since in the generate subtraction the activation related to the motor output associated with words has been subtracted away, and this activation is anatomically distinct from the foci found in the motor subtractions.

There are also two other foci of activation in the inferior cerebellum, one along the midline anteriorly (Focus 35, Table V) and one located more posteriorly and bilaterally (Foci 33, 34, Table V).

Auditory (Generate auditory words - Repeat Auditory Words) (GAW)

In the generate subtractions (GVW, GAW), as in the repeat subtractions (RVW, RAW), there is overlap in the areas activated across modalities. For generation of uses using auditory input, there are regions similar to those activated by generation of uses from visual input in inferior prefrontal cortex (Fig. 10, Focus 37; Table VI), anterior and inferior anterior cingulate (Foci 38, 39; Table VI), and anterior (Focus 11) and inferior lateral (Fig. 12, Focus 40, Table VI) cerebellum.

For the prefrontal (Foci 37) and inferior anterior cingulate (Focus 39) responses, there are anatomical dissimilarities with their counterpart foci

from GVW visual subtraction that may represent a modality specific but similar computation taking place in adjacent anatomical locations.

The small number of foci that are present for only one modality of presentation include the more superior prefrontal activations (foci 28,29; Table V), and the right posterior cerebellum (Focus 39, Table V) for GVW.

DISCUSSION

The results reported here represent our first attempt to extend the PET activation techniques of image subtraction and image averaging to "higher brain function", in particular to issues in lexical processing. The results have implications on several fronts. First, there is the information that can be obtained about how lexical information processing is represented in the brain. We believe our results agree quite well with multiple route models of lexical processing advanced by cognitive science (Coltheart, 1981; 1985; Marshall & Newcombe, 1973), and disagree substantially with the serial models (e.g. Geschwind, 1965, 1979). A second issue is the broader utility of PET activation studies in the neurobiology of human cognition and behavior. It seems reasonable that the techniques employed in these lexical studies could be extended to other cognitive questions as well. In order to assess the applicability of this technology to other human neurobiological and cognitive questions, questions about resolution, localization, and sensitivity of the techniques will be addressed. A third important topic is the applicability of the PET activation technology to clinical questions.

The representation of lexical processing in the brain

To assess the implications of our results for the representation of lexical processing in the brain, a two stage discussion of the results will be follow. First, we will focus on individual regions that are activated in our tasks and come up with reasonable hypotheses about the specific type of processing that might be taking place within that region. Next we will then attempt to arrange these functional regions into a framework that is consistent with our data and some of the other information extant on lexical processing.

Regions of activation

The regions in the occipital region activated in the passive visual words subtraction represent stages of visual processing. The activation of striate cortex is to be expected; the activation of the extrastriate regions is more interesting. The lack of activation in the extrastriate regions activated by visual words when other types of visual stimuli such as counterphase-flickered checkerboards, or dynamic random dot displays give preliminary evidence that this extrastriate activation is quite specific.

Some models of word processing, (LaBerge & Samuels, 1974, McClelland & Rumelhart, 1982) postulate levels of feature, letter, and word form analysis. In connectionist models (Rumelhart, 1986) these levels are reciprocally connected in a parallel distributed fashion. The existence of multiple areas of activation in the occipital lobe is consistent with such a multi-level processing network, with the end level representing activation at a visual

word form level of analysis. Further evidence that visual word form codes might be present in anterior occipital areas is that lesions of this area in the dominant hemisphere sometimes cause pure alexia, that is, the inability to read words (although letter by letter reading may still occur) without other language deficits (Damasio & Damasio, 1983; Henderson, 1986). Letter by letter reading has been attributed to a deficit in the visual word form (Warrington & Shallice, 1980) but could also involve earlier stages of the network (Shallice & Saffran, 1986). Our overall results are certainly consistent with an important role of the occipital temporal junction in processing words either passively or actively. However, the PET results do not reflect the degree of lateral asymmetry found in lesions.

In audition as in vision, there appears to be activation in the primary sensory receiving zone, and also in secondary regions (SMG, inferior temporal cortex) that are not activated by simple stimuli, such as clicks, tones, or noise bursts. It is tempting to hypothesize that these secondary regions of activation may represent a phonological level of encoding. Lesions of the supramarginal gyrus have been shown to produce a phonological agraphia in which patients can write words from dictation but not nonwords (Roeltgen, Sevush & Heilman, 1983; Shallice, 1981). This result has identified the left supramarginal gyrus with phonological as distinct from auditory processing. The activation of the left SMG during passive auditory word presentation (but in unpublished work from this laboratory not for auditory clicks or tones) makes the SMG a good candidate for phonological encoding of words at least from auditory input.

Several foci are activated when subjects repeat words presented either visually or auditorily. Activation in motor cortex and cerebellum would be expected in any motor task, and is present for verbal output. Several other areas are activated as well, areas that have been implicated in some aspect of motor coding or programming including the SMA, bilateral activation inferior premotor "Broca's" cortex, and a left lateralized premotor region superior to the Broca's activation.

While Broca's area has been implicated in language-specific deficits, a review by Mohr et al, (1978) has shown that lesions that are confined to this region produce only motor and praxis deficits, without specific language involvement, and that to produce the full-blown syndrome of Broca's aphasia requires larger lesions that extend more anteriorly. Our results showing that simple tongue movement also activates this region is consistent with this view. Recent studies showing that actual and imagined hand movement activate cortex at or near this site make it likely that the "Broca's" activation plays a role in higher level motor coding or production.

SMA is activated in conditions similar to that of our inferior premotor region. It is activated during verbal output and silent counting of associated pairs. This area has been hypothesized to act as a stage in programming of complex movements (Roland, Larson, Lassen & Skinhoj, 1980) or as a stage in a medial motor system related to lower motor process (Goldberg, 1985). The activations in our lexical processing tasks are not inconsistent with these hypotheses. However, SMA is also activated during simple hand movements, and during visual imagination of simple hand movements (Fox, et al, 1987) While all of the tasks include at least an implied output (silent

counting, verbalization, or imagined or actual hand movement), neither an actual movement, nor complex programming appears to be necessary to activate this region.

Potential explanations for activation in the generate task include semantic association, grammatical association, spontaneous generation from internal representation (since this task is not directly input driven), and possible non-specific arousal (since this task is the most difficult). The areas activated by the generate use stage include cingulate cortex, left inferior frontal cortex.

Some literature has implicated cingulate cortex with emotion, or autonomic response (Burns & Wyss, 1985), so it is possible that the activation there is related to the nonspecific arousal from task difficulty. A more intriguing possibility is that the ACC activation is related to spontaneous generation, that is, to the initiation or production of a response that does not represent a simple sensory to motor translation. Large midline lesions including SMA & ACC often produce akinetic mutism, a syndrome in which spontaneous speech is extremely rare (Masdau, Schoene, Funkenstein, 1978; Barris & Schuman, 1953; Nielsen & Jacobs, 1951). This deficit is consistent with ACC activation in the generate task, since our generate task demands a spontaneous associated response.

Mohr's (et al, 1978) study of inferior frontal lesions showed that large lesions that extended anteriorly to Broca's area were needed to produce full-blown Broca's aphasia. Studies by Goldman-Rakic (1987) have implicated similar areas in monkeys as being involved in higher order transformations or representations of information. An animal with lesions to this area has difficulty in withholding a preponent, direct response to the stimulus when the animal is asked to hold and transform the information and act on that transformed representation. Goldman also hypothesizes that different subregions of dorsolateral prefrontal cortex may perform similar transformations on different input information, and the adjacent but slightly different localizations for auditory and visual presentations would be consistent with this hypothesis. Since there is convergence of activation on two semantic association tasks, the inferior anterior prefrontal activation in the generate and semantic monitoring task is a candidate for computation related to semantic association.

The activation in right lateral inferior cerebellum is anatomically distinct from activation found with the repeat words and other motor tasks. The different response locale from cerebellar motor activation and the presence of the activation in the generate use subtractions argue for cognitive rather than sensory or motor computation being related to this activation. A role for the cerebellum in cognition has been advanced in recent papers (e.g. Leiner, Leiner & Dow, 1987), but there appears to be no specific candidate set of computations required by our generate task that might be related to cerebellar activation.

Models of lexical processing

The study of lexical processing has taken two relatively independent courses. There have been studies from a neurological orientation with a

focus of correlating specific brain regions with particular subcomponents of language processing (See Henderson 1986 for a recent review). These studies have tended to focus on observations and tests on individuals with brain injuries that affect aspects of their language processing or speech production. The most influential synthesis of these studies is in the disconnection models of Geschwind (1965, 1979). His interpretations of these results, while not substantially different from other accounts that had been offered in the late 1800s (See Henderson 1986) were so compelling that they have become the standard neurological model. This model is essentially serial in its analysis of aphasias, and related problems. For example, to read aloud a printed word, the visual information must first be processed in the visual cortex. This visual information is then sent to the angular gyrus where the visual word forms organize the letters, and transform the organized form to an auditory form. Wernicke's area holds a concomitant auditory word memory and directs associations so that an analysis of the word meaning takes place. This code is then sent to Broca's area, where the articulation of the word is prepared, and then to primary motor cortex for output. When this serial neurological model was advanced in the middle 1960's, serial models were in vogue in other areas of neuroscience as well. This was the era when the simple to complex to hypercomplex model of visual information processing (Hubel & Wiesel, 1962, 1965) was having a great impact. Also, in the absence of information to the contrary, a serial model would reasonably be considered a best first guess.

Over the past twenty years, cognitive studies of word processing have focused on different aspects of the problem (See Coltheart, 1985). Cognitive scientists have been more interested in the elementary cognitive operations that make up language than in the anatomical substrates underlying these operations (Mehler, Morton & Jusczyk, 1984). As such, their experiments have been directed toward dissociations of the different aspects of lexical processing. Many experiments take the form of reaction time studies of normals, but lesion behavior studies have been performed as well. In the lesion-behavior studies, the focus has also been to define mental operations that are dissociated from one another by brain injury rather than on associating the loss of a particular cognitive operation with a particular lesion. For example, can a brain-injured individual be found who can pronounce a familiar visually presented word, such as colonel, while being unable to translate a simple pronounceable nonword such as "caik" into a phonological representation that could then be said aloud? Such a dissociation between a visual word form, and phonological encoding of simple pronounceable visual nonword form can be shown to exist, and is strong evidence against a single pathway serial model (Coltheart, 1981; 1985, Humphreys & Evett, 1985).

Recent cognitive models for lexical processing have tended to focus on a variations of a dual route approach for reading of visual words (Coltheart, 1985; Humphreys & Evett, 1985; Carr & Pollatsek, 1985). In these models, visual information has relatively independent access to phonological information about the word (phonological codes), and information about the meaning of the word (semantic codes). In other words, separate phonological and semantic pathways are postulated to arise from visual information about the word.

Our first impetus for investigating the multiple route models were our early results from the PET visual word processing experiments. In the visual task conditions that we employed, there was never any evidence of activation in Wernicke's area or angular gyrus in any of the conditions. When reading a word aloud, the serial neurological model would have predicted activation in both of these regions (Geschwind, 1979). One potential explanation for this result could be that our PET activation techniques were not sensitive enough to see activation in these areas, perhaps because of a particularly high degree of anatomical variability in these regions that subverted the image averaging process. This explanation seems unlikely in that we find clear activation near this region during auditory word presentation (SMG activation). The other explanation is that a strictly serial model for word processing is insufficient. The evidence for independent pathways from cognitive literature argues that this is the case, and our results confirm that there is more than one pathway for the processing of visual words. We will now review some of the evidence for separate routes to word processing in light of cognitive studies, lesion studies, and our PET activation findings, and attempt to find a tentative framework that is consistent with many of these results.

To build up this framework, we will focus on several issues of dissociation that make it likely that there are separate processing routes for the different types of information. These dissociations include independent pathways for the generation of word codes for visual and phonological information, independence of phonological input and articulatory output codes, and independence of semantic association and articulatory codes. In this context, independence between codes is defined as the ability for information used to generate one code not obligatorily activating or generating the other code. We will assess this independence in several ways. From cognitive literature, we will use information from dual task performance and other tests that attempt to define independent processing modules. The logic of dual task interference studies is that when performing two tasks, the amount of deficit that the performance of the second task adds to the first is a measure of the dependence on shared pathways of information processing between the two tasks. From lesion-behavior studies, we will attempt to find behavioral dissociations such that a lesion affects one type of behavior, while leaving another type of behavior relatively intact. From our PET results we will assess the conditions that do or do not activate a region in an attempt to correlate an area with processing demands of the activity tasks.

Our first evidence from the PET studies came during the visual studies in the lack of activation in angular gyrus-Wernicke's region during any of the visual reading tasks. This made it seem reasonable that the visual information, perhaps in the form of a visual word code generated in the lateral occipital extrastriate cortex, had relatively direct access to articulatory output. This made it unlikely that a phonological input-type recoding of the word had to take place for output to occur.

There is considerable evidence on the independence of visual from phonological codes in the cognitive literature of visual word reading (See Carr & Pollatsek, 1985 for a review). According to these studies, visual word forms have direct access to semantic and articulatory codes using a

lexical route as well as access to phonological codes. Once source of evidence is the ability of subject to pronounce irregular words (e.g. pint and colonel) rapidly (Baron & Strawson, 1976). A second line of research shows that visual factors rather than phonological ones, influence the time to determine if a letter string is a lawful English word (Coltheart, Davelaar, Jonasson & Besner, 1977) and whether a word string (e.g. tie the not) is meaningful (Baron, 1973). Finally, dual tasks in which subjects are required to process visual words while also processing auditory information show that the secondary auditory tasks affect rhyming judgments but often do not affect meaning judgments (Kleiman, 1975; Rollins & Hendricks, 1980). These data have usually been interpreted as suggesting that in the skilled reader visual word forms have direct access to areas for articulation and semantics.

Studies of individuals with brain injuries also suggest the existence of somewhat separate routes. There are reported brain injury cases in which individuals correctly pronounce regular words and nonsense syllables but who mispronounce irregular words such as "pint" and "colonel" in accordance with the rules for English. These people have difficulty with the direct route from visual word form to articulation. Other patients correctly pronounce these irregular words, but have difficulty with constructions such as "caik", even though they sound like a real word that they can readily pronounce (Coltheart, 1981; 1985).

The PET results, the cognitive studies of normals and the studies of dissociations in dyslexia provide support for a direct route from visual word forms to semantic and articulatory codes.

There also seems to be independence in the phonological input and output codes. It has been postulated that a single representation of the sound of a word could suffice for decoding auditory input and for encoding word output. The PET results would argue against this. While there appears to be a common region activated for word output for both visual and auditory presentation near classically defined Broca's area, this area is not activated by auditory word input. Dual task studies by Shallice, McLeod & Lewis (1985) confirm the necessity of separate representations for phonological input and articulatory output. In these studies, a load was placed on the output system by having people read aloud visually presented words. When this was combined with a task where people were to monitor for a target word in a stream of words presented auditorily, there was very little dual task interference. This not only argues for separate speech input and output codes, but is also confirmatory evidence again that visual reading does not require (interfere with) phonological encoding. Many lesion studies as well confirm that auditory processing of words (comprehension) can be impaired while leaving articulation relatively intact, and articulation be affected while leaving processing of auditory words intact (Broca's vs. Wernicke's Aphasia, Geschwind, 1965).

It is evident that we can pronounce words or letter strings the meaning of which we do not know. Our PET results agree that it is possible to process and articulate words without measurable activation of regions related to semantic association. The inferior frontal area that is activated in semantic association tasks, is not activated during the repeat condition.

Lesions that produce a syndrome called isolation aphasia confirm that a pathway that allows pronunciation without semantic processing might exist. In isolation aphasia, repetition is relatively spared while comprehension is deficient. Often, a syndrome known as echolalia can result, in which the individual parrots whatever is heard without apparent comprehension. James (1890) addressed this point to the effect that "it is more difficult to ascend to the meaning of a word than to pass from one word to another; or to put it otherwise, it is harder to be a thinker than to be a rhetorician, and on the whole nothing is commoner than trains of words not understood."

Given these associations, a tentative cognitive-anatomical framework is presented in Fig. 3. This represents a best guess arrangement of cognitive operations and their associated anatomical region from the evidence described above. While we do not attempt a complete breakdown of the cognitive components in lexical processing, it provides a guiding framework for further experiments is provided.

Application of PET to studies of higher function

We believe that these and related studies using similar techniques can answer questions often raised about the use of PET for structure-function correlation. The questions involve matters of resolution, localization and sensitivity.

The resolution of the reconstructed images using PETT VI is 1.8 cm. The resolution is the distance between two simultaneously presented sources of radiation at which the two become resolvable as independent sources. More directly relevant to activation studies like those presented here is not the resolution of the image, but the localization of single point sources. In a study of the mapping of the retina onto striate cortex, changes in the location of sequentially imaged point sources of less than .5 cm. were consistently obtained (Fox, et al, 1986 Nature). Localization on this order seems appropriate for the studies that we have performed. The fact that we can find consistent localization across overlapping groups of subjects between modalities for regions such as rolandic motor mouth cortex or lateral cerebellum, confirms the reliability of the localization techniques that we employ.

The question of sensitivity presents a more complex problem. Our results indicate that consistent activation attributable to strictly cognitive operations is obtainable; in other words, our techniques have a level of sensitivity that produces consistent results. Both in the generate - repeat, and the converging covert monitoring tasks, areas are active that are very likely related to some level of semantic processing. The use of convergent paradigms to assess the association of an elementary cognitive operation with a particular anatomical region seems to us to be a fruitful way to proceed with the study of localization of function using PET activation techniques. On the other hand, studies from cognitive psychology argue that activation of associated words occur automatically even when the subject is unable to identify the word (Marcel, 1987). Priming studies show that presentation of an associate (e.g. doctor) will speed pronunciation or word-nonword judgments for a related word (e.g. nurse). In our passive conditions we see no activation that is easily attributable to semantic level coding, yet with 150

msec unmasked presentation there is little doubt that semantic priming would occur. The best interpretation would suggest that our PET method in measuring the activation of areas related to higher function is sensitive to information that is coded and used by the person in deliberately performing the task. Representation passively activated by input but not used in the execution of the task may be below the threshold of our PET method.

Another important issue in the application of our PET methodology involves the subtractive assumptions. Cognitive psychologists often object to the subtraction of two different tasks on the grounds that subjects may adopt entirely different strategies for the two tasks. Our PET methodology provides a means of checking up on this idea. Consider the comparison called generate - repeat. If generate involves all the processes involved in repeat plus some additional ones, subtracting fixation from generate should produce a pattern of activation like that found in the passive - fixation condition plus repeat - passive condition plus the additional areas in the generate - repeat condition. In fact, when we make such a subtraction, we find activation in several additional areas not found in generate - repeat. These include the cerebellum, SMA and areas around and Broca's area, and SMG and auditory receiving areas (Table VII). To a first approximation, our subtractive assumptions appear to be true for this set of tasks.

Clinical Implications

Can the view outlined in Figure 13 be reconciled with the lesion data that has supported the Geschwind model? For example, it is known that lesions of the angular gyrus produce alexia with agraphia. Our PET method only shows the activation of nuclear areas and not of white matter tracts. There has been a long unresolved discussion (see Henderson, 1986) in clinical neurology whether angular gyrus lesions have effects on the cortex or on fibers passing under the gyrus. If the routes from the visual system to frontal areas pass under the gyrus one would expect alexias if these fiber tracts are compromised. In addition, many readers do use phonological codes for individual words and readers generally require such codes in storing information during processing of passages.

Another discrepancy between our results and many clinical views is in the location of semantic association. It is well known that patients with lesions in Wernicke's area produce speech which is fluent but empty of semantic content. This suggests that Wernicke's area is involved in semantic processing. Recent evidence from studies of semantic priming has suggested however, that patients with anterior lesions and not posterior lesions have difficulty in access to semantic codes of individual items (Milberg & Blumstein, 1981; Milberg, Blumstein & Dwoetzy, 1987). It seems quite possible that the problems of Wernicke's aphasics involve the integration of semantic codes during the processing of word strings, possibly using phonological information, a level of processing not addressed in our tasks.

The current direct clinical utility of PET activation techniques is limited, but the ability to determine the anatomical location of areas related to specified functions could have clinical implications in the future. For example, our findings of lateralized foci in certain of the lexical processing tasks could be the basis for a method for determining

language dominance that is less invasive than the Wada hemispheric dominance test. Preliminary tests using these tasks to assess language dominance have been encouraging (Pardo, Fox, Goldring, Raichle, 1987).

In general, PET activation studies can be used to confirm and extend the catalogue of anatomical-functional relationships. This information can then be used in behavioral neurological diagnosis and would allow neurosurgeons to operate with greater confidence knowing the precise location of vital areas in relation to various surgical procedures.

General Discussion

Rather than an endpoint, we view these studies as a preliminary example of the utility of combining state of the art PET techniques with paradigms designed to address specific cognitive questions. The results lead us to tantalizing hypotheses about the representation of different aspects of lexical processing in the brain. Further experiments can be designed that directly address these hypotheses. For example, if a person must make a phonological (rhyme) judgment about visual word input, is our candidate region for phonological word coding, the supramarginal gyrus, activated?

Successful experiments along these lines could generate significant information about the neural mechanisms related to the processing of single words.

The methods employed here also represent a set of techniques that can be used in the study of the neural mechanisms underlying other higher functions such as attention, perception, and motor control. The results from PET activation studies can be used to complement and extend results using other techniques in both humans and other animals. As a result, the techniques can function in an integrated approach to basic neurobiological questions about higher processes.

REFERENCES

1. Barris, R.W. & Schuman, H.R. Bilateral anterior cingulate gyrus lesions. Neurology (Minneap.), 3 (1953) 44-52.
2. Baron, J. (1973) Phonemic stage not necessary for reading. Quart. J. of Exp. Psych. 25, 241-246.
3. Baron, J. & Strawson, C. (1976) Use of Orthographic and word specific knowledge in reading words aloud. J. of Exp. Psych. Human Perception and Performance 2, 386-393.
4. Burns, S.M. & Wyss, J.M. (1985) Involvement of the anterior cingulate cortex in blood pressure control. Brain Research, 340:71-77.
5. Carr, T.H. & Pollatsek, A. Models of word recognition in D. Besner, T.G. Wallker & G.E. McKinnon (eds.) Reading Research Vol. 5, 1985, Academic Press:New York 1-82.
6. Coltheart, M. Disorders of reading and their implications for models of normal reading. Visible Language, 1981, 15, 245-286.
7. Coltheart, M. (1985) Cognitive neuropsychology and the study of reading. IN M.I. Posner and O.S.M. Marin (eds.), Attention and Performance XI, Lawrence Erlbaum Associates; Hilldale, N.J.
8. Coltheart, M., Davelaar, E., Jonasson, J. & Besner, D. (1977) Access to the internal lexicon. In S. Dornic (ed.) Attention and Performance VI New York:Academic Press.
9. Damasio, A.R. & Damasio, H. The anatomic basis of pure alexia, Neurology, 1983, 33, 1573-1583.
10. Fox, P.T., Burton, H. and Raichle, M.E. (1987) Mapping human somatosensory cortex with positron emission tomography. J. Neurosurg., in press.
11. Fox, P.T., Mintun, M.E., Raichle, M.E., Miezin, F.M., Allman, J.M. and Van Essen, D.C. (1986) Response localization can surpass image resolution: Mapping the human visual system with positron emission tomography. Nature, 323: 806-809.
12. Fox, P.T., Miezin, F.M., Allman, J.M., Van Essen, D.C. and Raichle, M.E. (1987a) Retinotopic organization of human visual cortex mapped with positron-emission tomography. J. of Neurosci. 7(3):913-922.
13. Fox, P.T., Pardo, J.V., Petersen, S.E. and Raichle, M.E. (1987b) Supplementary motor and premotor responses to actual and imagined hand movements using positron emission tomography. Soc. Neurosci. Abstr. 13: .
14. Fox, P.T., Perlmutter, J.S., and Raichle, M.E. (1985) A stereotactic method of anatomical localization for positron emission tomography. J. Comp. Assist. Tomogr. 9:141-153.

15. Fox, P.T. and Raichle, M.E. (1986b) Focal physiological uncoupling of cerebral blood flow and oxidative metabolism during somatosensory stimulation in human subjects. Proc. Natl. Acad. Sci. 83: 1140-1144.
16. Fox, P.T., Raichle, M.E., and Thach, W.T. (1985) Functional mapping of the human cerebellum with positron emission tomography. Proc. Natl. Acad. Sci., 82:7462-7466.
17. Geschwind, N. & Levitsky, W. (1968) Human Brain: left right asymmetries in temporal specific region. Science, 161:186-187.
18. Geschwind, N. (1965) Disconnection syndromes in animals and man. Part I-II. Brain, 88:237-294, 585-644.
19. Geschwind, N. Specialization of the human brain. Scientific American, 1979, 158-168.
20. Gibson, E.J., Shurcliff, A., and Yonas, A. (1970) Utilization of spelling patterns by deaf and hearing subjects. In Basic Studies in Reading, H. Levin and J. Williams (eds.), New York: Basic Books.
21. Goldberg, G. (1985) Supplementary motor area structure and function: Review and hypotheses. The Behavioral and Brain Sciences, 8:567-616.
22. Goldman-Rakic, P.S. (1987) Circuitry of primate prefrontal cortex and regulation of behavior by representational knowledge. In: Higher Cortical Function; (F. Plum and V. Mountcastle, Eds.) Handbook of Physiology, Amer. Physiol. Soc., Washington, D.C., 5:373-417.
23. Henderson, V.W. Anatomy of posterior pathways in reading: a reassessment. Brain and Language (1986) 29, 199-133.
24. Humphreys, G.W., and Evett, E.J. (1985) Are there independent lexical and nonlexical routes in word processing? An evaluation of the dual route theory in reading. The Behavioral and Brain Sciences, 8:689-740.
25. Imig, T.J., Ruggero, M.A., Kitzes, L.M., Javel, E. and Brugge, J.F. Organization of auditory cortex in the owl monkey (*Aotus trivirgatus*). J. Comp. Neurol. 171:111-128.
26. James, W. Principles of psychology. (1980) New York:Holt.
27. Kleiman, G.M. (1975) Speech recoding in reading. Journal of Verbal Learning and Verbal Behavior, 24, 323-339.
28. LaBerge, D. & Samuels, J. Toward a theory of automatic information processing in reading. Cognitive Psychology, 1974, 6, 293-323.
29. Leiner, H.C., Leiner, A.L. & Dow, R.S. (1986) Does the cerebellum contribute to cognitive skills? Behavioral Neuroscience, August.
30. Marcel, A.J. Conscious and unconscious processes in word recognition. Cognitive Psychology, 1983, 15, 197-237.

31. Marshall, J.C. & Newcombe, F. Patterns of paralexia: a psycholinguistic approach. J. of Psycholinguistic Research, 1973, 2, 175-198.
32. Marslen-Wilson, D.W. Functional parallelism in spoken word-recognition. Cognition, 1987, 25, 71-102.
33. Masdau, J.C., Schoene, W.C. & Funkenstein, H. (1978) Aphasia following infarction of the left supplementary motor area. Neurology, 28:1220-23.
34. McCarthy, R. & Warrington, E.K. Category specificity in an agrammatic patient: the relative impairment of verb retrieval and comprehension. Neuropsychologia, 1985, 23, 709-727.
35. McClelland, J.L. & Rumelhart, D.E. (1981) An interactive activation model of context effects in letter perception: Part 1. An account of basic findings. Psychological Review, 88:375-407.
36. Mehler, J., Morton, J. & Jusczyk, P.W. On reducing language to biology. Cognitive Neuropsychology, 1984, 1, 83-116.
37. Milberg, W., Blumstein, S.E. & Dworetzky, B. Processing of lexical ambiguities in aphasia. (1987) Brain and Language, 31, 138-150.
38. Milberg, W. & Blumstein, S. (1981) Lexical decision and aphasia: evidence for semantic processing. Brain and Language, 14, 371-385.
39. Mintun, M.A., Fox, P.T. and Raichle, M.E. (1987) Signal averaging applied to positron emission tomography. Soc. for Neurosci. Abstr. 13: .
40. Mohr, J.P., Pessin, M.S., Finkelstein, S., Funkenstein, H.H., Duncan, G.W. and Davis, K.R. (1978) Broca aphasia: Pathologic and clinical. Neurology, 28:311-324.
41. Nielsen, J.M. & Jacobs, L.L. Bilateral lesions of the anterior cingulate gyri. Bull. Los Angeles Neurol. Soc. 16 (1951) 231-234.
42. Perlmuter, J.S., Herscovitch, P., Powers, W.P., Fox, P.T., and Raichle, M.E. (1985) A standardized mean regional method for calculating global positron emission tomographic measurements. J. Cereb. Blood Flow and Metab. 5:476-480.
43. Petersen, S.E., Fox, P.T., Posner, M.I., Mintun, M.A., and Raichle, M.E. (1986) Focal brain activity during visual language tasks as measured with averaged PET images of evoked CBF change. Soc. Neurosci. Abstr., 12:1161.
44. Posner, M.I. (1978) Chronometric explorations of mind. Englewood Heights N.J.:Lawrence Erlbaum Assoc.
45. Roeltgen, D.P., Sevush, S. & Heilman, K.M. Phonological agraphia, Neurology, 1983, 33, 755-765.

46. Roland, P. Cortical organization of voluntary behavior in man. Human Neurobiology, 1985, 4, 155-167.
47. Roland, P., Larson, B., Lassen, N.A. & Skinhoj, E. (1980) Supplementary motor area and other areas in organization of voluntary movements in man. Journal of Neurophysiology. 43, 118-136.
48. Rollins, H.A. & Hendricks, R. (1980) Processing of words presented simultaneously to eye and ear. Journal of Exp. Psych:Human Perception and Performance, 6, 99-109.
49. Rumelhart, D.E. and McClelland, J.L. (1982) An interactive activation model of context effects in letter perception: Part 2. The contextual enhancement effect and some tests and extensions of the model. Psychological Review 89: 60-94.
50. Rumelhart, D.E. & McClelland, J.L. (1986) Parallel distributed processing. Vol. 1 and 2., MIT Press:Cambridge.
51. Shallice, T. (1981) Phonological aphasia and the lexical route in writing. Brain, 104, 413-429.
52. Shallice, T., McLeod, P. & Lewis, K. Isolating cognitive modules with the dual task paradigm: are speech perception and production separate processes? Quarterly Journal of Experimental Psychology, 1985, 37A, 507-532.
53. Shallice, T. & Saffaran, E. Lexical processing in the absence of explicit word identification. Cognitive Neuropsychology, 1986, 3(4), 429-458.
54. Snedecor, G.W. & Corcoran, W.G. (1980) Statistical Methods, Iowa University Press.
55. Sternberg, S. The discovery of processing stages: Extensions of Donders' method. Acta Psychologica, 1969, 30, 276-315.
56. Warrington, E.T. & Shallice, T. Word form dyslexia, Brain, 1980, 103, 99-112.

Table legend

For all of the tables in this paper the following conventions are used: the region is given a mnemonic anatomical name associated with the coordinates. The coordinates and magnitudes of response are determined using a three dimensional search algorithm on the averaged subtraction image. The coordinates are in mm. from a 0,0,0 point that is at the level of a line drawn between the anterior and posterior commissures ($z=0$), at the midline of the brain ($x=0$), and located anterior posteriorly halfway between the commissures ($y=0$). The magnitudes are the change in blood flow in gm/100ml/min, and the statistical significance of the points are assessed with a two stage testing procedure. The distribution of the magnitudes of local BF change is tested for outliers using an omnibus gamma2 test. For all averaged images presented here, there are statistically significant outliers. The foci with the largest magnitude of BF change are then given a z-score with respect to the population of all local changes within an image. All foci of change with a p-value $< .03$ are reported in the tables. $*--p < .03$, $**--p < .01$.

There is also a column in which other subtractions with anatomically similar significant foci of change are listed. $+-$ there is a region in a location near to the activation focus listed, but may be anatomically distinct.

For table 7, the 15 foci with the greatest magnitude of change are reported irrespective of their statistical significance since this table is for comparison purposes. $\#--p > .03$

In general, the passive presentation subtractions identify modality specific foci of activation, while the higher level subtraction generally activate similar regions across modalities. Table VII shows that to a first approximation our subtractive assumptions are correct, since this cross level subtraction contains regions which are the sum of the three subtraction levels within it (see discussion).

Table I

Subtraction Conditions Passive Visual Words - Fixation Point (PVW)N = 7; omnibus test, $p < .01$

Region		COORDINATES (mm)			Magnitude	Other conditions w/similar activation foci
		Z	X	Y		
1. Striate Cortex	(L)	10	6	-72	2.28*	---
2.	(R)	10	-12	-72	2.66*	---
3. Extrastriate Cortex	(L)	2	24	-58	3.82**	---
4.	(R)	6	-26	-66	2.95**	---
5. Inferior Lateral Occipital Cortex	(R)	-4	-34	-46	3.38**	---
6. Putamen	(L)	4	22	24	3.32**	---

Table II

Subtraction Conditions Passive Auditory Words - Fixation Point (PAW)N = 8; omnibus test, $p < .05$

Region	COORDINATES (mm)			Magnitude	Other conditions w/similar activation foci
	Z	X	Y		
7. Posterior Superior Temporal Cortex (L)	14	46	-10	2.46*	---
8. (R)	12	-42	-16	2.76**	---
9. Inferior Temporal Cortex (L)	-2	42	10	3.02**	---
10. Supramarginal Gyrus (L)	14	54	-30	2.88**	---
11. Lateral "Broca" Cortex (R)	8	-62	-12	3.30**	RVW, RAW
12. Inferior Anterior Cingulate Cortex (L)	18	12	44	2.34*	---

Table III

Subtraction Conditions Repeat Visual Words - Passive Visual Words (RVW)N = 11; omnibus test, $p < .01$

Region		COORDINATES (mm)			Magnitude	Other conditions w/similar activation foc
		Z	X	Y		
13. Mouth Region,						
Rolandic Cortex	(L)	40	46	0	4.34**	RAW
14.	(R)	32	-52	6	3.46**	RAW
15. Buried "Broca"						
Cortex	(L)	14	31	6	3.04*	RAW
16. Lateral "Broca"						
Cortex	(R)	8	-63	-4	2.96*	RAW, PAW
17. Premotor Cortex	(L)	18	48	14	2.98*	RAW ⁺
18. SMA		50	-2	10	3.36*	RAW
19. Superior Anterior						
Cerebellum	(L)	-8	6	-42	4.62**	---
20.	(R)	-9	-16	-44	4.47**	---
21. Colliculus		-6	0	-24	3.04	GAW

Table IV

Subtraction Conditions Repeat Auditory Words - Passive Auditory Words (RAW)N = 10; omnibus test, $p < .05$

Region	COORDINATES (mm)			Magnitude	Other conditions w/similar activation for
	Z	X	Y		
22. Mouth Region, Rolandic Cortex (L)	42	46	-2	3.64**	RVW
23. (R)	40	-56	2	3.78**	RVW
24. Buried "Broca" Cortex (L)	14	34	10	3.17*	RVW
25. Lateral "Broca" Cortex (R)	12	-62	-7	3.22**	RVW, PAW
26. Premotor Cortex (L)	26	52	2	3.06*	RVW+
27. SMA	52	2	14	2.80*	RVW

Table V

Subtraction Conditions Generate Visual Words - Repeat Visual Words (GVW)

N = 12, p < .01

Region	COORDINATES (mm)			Magnitude	Other conditions w/similar activation for
	Z	X	Y		
28. Dorsolateral Pre- frontal Cortex (L)	20	44	36	2.98**	GAW ⁺
29. Lateral Prefrontal Cortex (L)	8	38	36	2.96**	GAW ⁺
30. Inferior Prefrontal Cortex (L)	-6	-28	50	2.26*	GAW ⁺
31. Anterior Cingulate	38	-6	24	3.12**	GAW
32. Inferior Anterior Cingulate	28	-2	34	2.76**	GAW ⁺
33. Posterior Cerebellum (L)	-22	16	-64	2.26*	---
34. (R)	-18	-10	-58	2.62**	---
35. Anterior Cerebellum/ Colliculus (L)	-16	0	-36	2.70**	GAW
36. Inferior Lateral Cerebellum (R)	-22	-38	-48	3.34**	GAW

Table VI

Subtraction Conditions Generate Auditory Words - Repeat Auditory Words (GAW)N = 7; omnibus test, $p < .01$

Region	COORDINATES (mm)			Magnitude	Other conditions w/similar activation for
	Z	X	Y		
37. Inferior Prefrontal Cortex (L)	-6	33	43	3.10**	GVW ⁺
38. Anterior Cingulate	38	7	28	3.28**	GVW
39. Inferior Anterior Cingulate	28	11	31	3.04**	GVW ⁺
40. Inferior Lateral Cerebellum (R)	-22	-37	-47	2.52**	GVW
41. Anterior Cerebellum/ Colliculus	-6	2	-30	3.00**	GVW

Table VII

Subtraction Conditions Generate Auditory Words - Fixation Point

N = 8, p < .01

Region	COORDINATES (mm)			Magnitude	Other conditions w/similar activation foc
	Z	X	Y		
Posterior Superior Temporal Cortex (L)	14	42	-16	2.76*	PAW
Supramarginal Gyrus (L)	14	54	-30	2.32#	PAW
Lateral "Broca" Cortex (R)	11	-63	-18	3.60**	PAW, RAW, RVW
Inferior Anterior Cingulate Cortex (L)	22	14	48	2.40#	PAW
Mouth Region, Rolandic Cortex (L)	37	47	4	3.02**	RAW, RVW
	32	-54	4	2.52#	RAW, RVW
Buried "Broca" Cortex (L)	14	42	10	4.44**	RAW, RVW
SMA	45	-2	16	4.66**	RAW, RVW
Superior Anterior Cerebellum (L)	-12	22	-42	4.04**	RVW
	-12	-12	-34	3.04**	RVW
Anterior Cerebellum/ Colliculus	-16	0	-30	3.40**	GAW, GVW
Anterior Cingulate	38	0	19	4.66**	GAW, GVW
Dorsolateral Prefrontal Cortex (L)	27	32	46	2.20#	GAW, GVW ⁺
Lateral Prefrontal Cortex (L)	16	48	29	2.58#	GAW ⁺ , GVW
Inferior Prefrontal Cortex	2	42	38	2.20#	GAW, GVW ⁺
Inferior Lateral Cerebellum	-19	-26	-44	3.22**	GAW, GVW
Putamen	4	23	27	2.70#	

- Figure 1. A. Primary image of brain blood flow (BF) while viewing fixation point only (control state). Each of the seven slices is a horizontal slice from top (1) to bottom (7), with anterior at the top and left left on the slice representing the left brain. B. Primary BF image acquired during peripheral field stimulation (5.5 to 15.5 degrees eccentricity). C. Subtraction image of the change in BF induced by peripheral visual stimulation, created by subtracting image A from image B. Peak response is on slices 3 and 4. All images are from a single subject.
- Figure 2. Schematic lateral surface view of the brain. Anterior is to the right. The horizontal lines mark two reference planes of section in the standardized stereotactic space. The 0.0 line runs through the anterior and posterior commissures (AC-PC line). The 4.2 line is parallel to the AC-PC line and 4.2 cm above it.
- Figure 3. Auditory vs. visual comparison. A horizontal slice through averaged subtraction image represents BF change when BF fixation is subtracted from BF present during presentation of word stimuli at 1 Hz (sensory subtraction). Slice is taken .2 cm below AC-PC line. Foci of activity present at this level includes inferior temporal non-primary auditory cortex in the left hemisphere (A) for auditorily-presented words, and extrastriate inferior occipital responses (B) for visually-presented words.
- Figure 4. Same condition as Fig. 3. Slice in Fig. 4 is taken 1.6 cm above AC-PC line. Foci of activity present at this level include supramarginal gyrus (SMG) (A), bilateral superior posterior temporal cortex (B), and inferior anterior cingulate (C) for auditory presentation, and some occipital cortical activation (D) for visual presentation. Note the non overlapping distributions of activity in Figs. 3 and 4 during passive presentation.
- Figure 5. Auditory vs. visual comparison. A horizontal slice through an averaged subtraction image representing blood flow (BF) change when blood flow during passive presentation of words is subtracted from blood flow during vocal repetition of presented words (motor subtraction). Slice is taken 4.0 cm. above AC-PC line. The foci present for both auditory and visual presentation are located on rolandic cortex, just anterior and superior to regions activated by somatosensory stimulation of the lips (Fox, , 1987) and likely represent the mouth representation of primary motor cortex.
- Figure 6. Same conditions as Fig. 5. Slice is taken 1.2 cm. above AC-PC line. Foci for both auditory and visual presentation are located on the frontal cortex (F3) buried in the sylvian sulcus. The left site of activation is similar in location to Broca's area, but our activation is clearly bilateral.

- Figure 7. Comparison of activation for repetition of words and simple tongue movement. The repeat words slice is taken from Fig. 6 (visual presentation of words); the tongue movement slice represents a subtraction of blood flow during an eyes closed resting state from simple side-to-side movements of the tongue throughout the scan. Similar activation is present in the two conditions, and also for actual and imagined hand movement (Fox, Pardo, ,1987).
- Figure 8. Same conditions as Figure 5. Foci for both auditory and visual representations occur near the midline on frontal cortex (F1). This region is commonly referred to as the supplementary motor area (SMA). This area appears to be active for all tasks that require motor programming (Fox, Pardo, 1987, Roland, , Goldberg, 1985).
- Figure 9. Auditory vs. visual comparison. A horizontal slice through an averaged subtraction image representing BF change when BF during repetition of presented words is subtracted from BF during vocalization of an appropriate use for the presented word (e.g. presentation of "cake"...output might be "eat") (cognitive subtraction). Slice is taken 3.4 cm. above AC-PC line. Foci for both auditory and visual presentation are in the anterior cingulate cortex.
- Figure 10. Conditions same as Fig. 9. Slice is taken .8 cm. below AC-PC line. Foci for both presentation modalities occur in inferior anterior frontal cortex, probably area 47 of Brodmann. Those areas of activation are strongly left lateralized.
- Figure 11. Comparison of activation in two semantic tasks. The slice on the right is the same as Fig. 9, visual presentation; the slice on the left represents the BF change when the BF during passive presentation of words at 2.5 Hz is subtracted from BF during a condition where the subject is asked to monitor this string of words for members of a specific semantic category. In the semantic monitoring task, there is no motor output during the scan. Subjects are asked after the scan for a gross estimate of the percentage of target words. The similar foci of activation in these two different semantic tasks implicate this region in some level of semantic processing.
- Figure 12. Conditions the same as Fig. 9. Slice taken 2.2 cm. below AC-PC line. Foci in both slices are strongly lateralized to the right and are located in the lateral cerebellum. The right lateralization (and its implied relationship to the left cerebral hemisphere) implicate the cerebellum in some higher cognitive function.
- Figure 13. A general network relating some of the areas of activation in this study to the different levels of lexical processing. There are many alternative networks consistent with the conditions under which the areas are activated, but this arrangement represents a simple design consistent with our results, and some convergent experiments from other types of studies.




FIGURE 1

- A. Primary image of cerebral blood flow (CBF) while viewing fixation point only (control state).
- B. Primary CBF image acquired during peripheral field visual stimulation (5.5 to 15.5 degrees eccentricity)
- C. Subtraction image of the absolute change in CBF induced by peripheral stimulation, created by subtracting image A from image B. Peak response is on slices 3 and 4. All images are from scanning a single subject.

FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12





FIGURE 13



(Washington University/Posner) 1987/11/10

Distribution List

Distribution List

Dr. Phillip L. Ackerman
University of Minnesota
Department of Psychology
Minneapolis, MN 55455

Dr. Beth Adelson
Department of Computer Science
Tufts University
Medford, MA 02155

Technical Director,
Army Human Engineering Lab
Aberdeen Proving Ground
MD 21005

Dr. Robert Ahlers
Code M711
Human Factors Laboratory
Naval Training Systems Center
Orlando, FL 32813

Dr. Ed Aiken
Navy Personnel RAD Center
San Diego, CA 92152-6800

Dr. John Allen
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Dr. Earl A. Alluisi
HQ, AFHRL (AFSC)
Brooks AFB, TX 78235

Dr. James Anderson
Brown University
Center for Neural Science
Providence, RI 02912

Dr. Nancy S. Anderson
Department of Psychology
University of Maryland
College Park, MD 20742

Technical Director, ARI
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Gary Aston-Jones
Department of Biology
New York University
1009 Main Bldg
Washington Square
New York, NY 10003

Dr. Alan Baddeley
Medical Research Council
Applied Psychology Unit
15 Chaucer Road
Cambridge CB2 2EF
ENGLAND

Dr. James Ballas
Georgetown University
Department of Psychology
Washington, DC 20057

Dr. Harold Bamford
National Science Foundation
1800 G Street, N.W.
Washington, DC 20550

Dr. Isaac Bejar
Educational Testing Service
Princeton, NJ 08540

Dr. Alvin Blittner
Naval Biodynamics Laboratory
New Orleans, LA 70189

Dr. John Blaha
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Dr. Sue Bogner
Army Research Institute
ATTN: PERI-SF
5001 Eisenhower Avenue
Alexandria, VA 22333-5600

Dr. Gordon H. Bower
Department of Psychology
Stanford University
Stanford, CA 94306

Mr. Donald C. Burg
General Physics Corp.
10650 Hickory Ridge Rd.
Columbia, MD 21044

Dr. Gail Carpenter
Northeastern University
Department of Mathematics, 504LA
360 Huntington Avenue
Boston, MA 02115

Dr. Pat Carpenter
Carnegie-Mellon University
Department of Psychology
Pittsburgh, PA 15213

Dr. Tyrone Cashman
American Society of
Cybernetics
3428 Fremont Ave. South
Minneapolis MN 55408

Dr. Alphonse Chapanis
8415 Bellona Lane
Suite 210
Buiton Towers
Baltimore, MD 21204

Dr. Paul R. Chatelier
OUSD&E
Pentagon
Washington, DC 20350-2000

Mr. Raymond E. Christal
AFHRL/HOE
Brooks AFB, TX 78235

Dr. David E. Clement
Department of Psychology
University of South Carolina
Columbia, SC 29208

Dr. Charles Clifton
Tobin Hall
Department of Psychology
University of
Massachusetts
Amherst, MA 01003

Assistant Chief of Staff
for Research, Development,
Test, and Evaluation
Naval Education and
Training Command (N-5)
NAS Pensacola, FL 32508

Dr. Michael Coles
University of Illinois
Department of Psychology
Champaign, IL 61820

Dr. Allan M. Collins
Bolt Beranek & Newman, Inc.
50 Moulton Street
Cambridge, MA 02138

Dr. Stanley Collyer
Office of Naval Technology
Code 222
800 M. Quincy Street
Arlington, VA 22217-5000

Dr. Leon Cooper
Brown University
Center for Neural Science
Providence, RI 02912

Dr. Lynn A. Cooper
Learning RAD Center
University of Pittsburgh
3939 O'Hara Street
Pittsburgh, PA 15213

Phil Cunliff
Commanding Officer, Code 7522
Naval Undersea Warfare Engineering
Keyport, WA 98345

Brian Dailman
3400 TTM/TTCIS
Lowry AFB, CO 80230-5000

LT John Deaton
OMR Code 125
800 M. Quincy Street
Arlington, VA 22217-5000

Dr. Stanley Deutsch
Committee on Human Factors
National Academy of Sciences
2101 Constitution Ave.
Washington, DC 20548

Dr. R. K. Disbakes
Associate Director for Life Sciences
AFOSR
Building AFB
Washington, DC 20332

(Washington University/Posner) 1987/11/10)

Distribution List

Dr. Daniel Copher
Industrial Engineering
& Management
TECHNION
Haifa 32000
ISRAEL

Dr. Sherrie Gott
AFHRL/WDJ
Brooks AFB, TX 78235

Jordan Graffman, Ph.D.
2021 Lyttonsaville Road
Silver Spring, MD 20910

Dr. Richard H. Granger
Department of Computer Science
University of California, Irvine
Irvine, CA 92717

Dr. Steven Grant
Department of Biology
New York University
1009 Main Bldg
Washington Square
New York, NY 10003

Dr. Wayne Gray
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Bert Green
Johns Hopkins University
Department of Psychology
Charles & 38th Street
Baltimore, MD 21218

Dr. James G. Greene
University of California
Berkeley, CA 94720

Dr. William Greenough
University of Illinois
Department of Psychology
Champaign, IL 61820

Dr. Stephen Grossberg
Center for Adaptive Systems
Room 244
111 Cunningham Street
Boston University
Boston, MA 02215

Dr. Muhammad K. Habib
University of North Carolina
Department of Biostatistics
Chapel Hill, NC 27516

Prof. Edward Haertel
School of Education
Stanford University
Stanford, CA 94305

Dr. Henry M. Halff
Halff Resources, Inc.
4918 33rd Road, North
Arlington, VA 22207

Dr. Nancy F. Halff
Halff Resources, Inc.
4918 33rd Road, North
Arlington, VA 22207

Dr. Ronald K. Hambleton
Prof. of Education & Psychology
University of Massachusetts
at Amherst
Hills House
Amherst, MA 01003

Dr. Cheryl Hamel
NTSC
Orlando, FL 32813

Mr. William Hartung
PEAM Product Manager
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Harold Hawkins
Office of Naval Research
Code 11K2PT
800 M. Quincey Street
Arlington, VA 22217-5000

Prof. John R. Hayes
Carnegie-Mellon University
Department of Psychology
Schenley Park
Pittsburgh, PA 15213

Dr. Joan I. Heller
505 Haddon Road
Oakland, CA 94606

Distribution List

Dr. Stephanie Doss
Code 6021
Naval Air Development Center
Warminster, PA 18974-5000

Dr. Emanuel Dornhein
University of Illinois
Department of Psychology
Champaign, IL 61820

Mr. Ralph Dusek
ARD Corporation
5457 Twins Knolls Road
Suite 400
Columbia, MD 21045

Dr. Ford Ebner
Brown University
Anatomy Department
Medical School
Providence, RI 02912

Dr. Jeffrey Eisan
University of California,
San Diego
Department of Linguistics, C-008
La Jolla, CA 92093

Dr. William Epstein
University of Wisconsin
W. J. Brogden Psychology Bldg.
1202 W. Johnson Street
Madison, WI 53706

Dr. K. Anders Ericsson
University of Colorado
Department of Psychology
Boulder, CO 80309

Dr. Jerome A. Feldman
University of Rochester
Computer Science Department
Rochester, NY 14627

Dr. Paul Feltoovich
Southern Illinois University
School of Medicine
Medical Education Department
P.O. Box 3926
Springfield, IL 62708

Dr. Craig I. Fields
ABPA
1400 Wilson Blvd.
Arlington, VA 22209

Dr. Gail R. Fleischaker
Margulis Lab
Biological Sci. Center
2 Cunningham Street
Boston, MA 02215

Dr. Jane M. Flinn
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Dr. Michaela Gallagher
University of North Carolina
Department of Psychology
Chapel Hill, NC 27514

Dr. R. Edward Geiselman
Department of Psychology
University of California
Los Angeles, CA 90024

Dr. Don Gentner
Center for Human
Information Processing
University of California
La Jolla, CA 92093

Dr. Lee Giles
AFOSR
Bolling AFB
Washington, DC 20332

Dr. Eugene E. Gloye
Office of Naval Research
Detachment
1030 E. Green Street
Pasadena, CA 91106-2485

Dr. Joseph Goguen
Computer Science Laboratory
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025

(Washington University/Posner) 1987/11/10

Distribution List

Dr. Per Heimerlsen
University of Oslo
Department of Psychology
Box 1098
Oslo 3, NORWAY

Dr. Steven A. Hillyard
Department of Neurosciences
University of California,
San Diego
La Jolla, CA 92093

Dr. Geoffrey Hinton
Carnegie-Mellon University
Computer Science Department
Pittsburgh, PA 15213

Dr. Jim Hollan
Intelligent Systems Group
Institute for
Cognitive Science (C-015)
UCSD
La Jolla, CA 92093

Dr. John Holland
University of Michigan
2313 East Engineering
Ann Arbor, MI 48109

Dr. Melissa Holland
Army Research Institute for the
Behavioral and Social Sciences
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Keith Holyoak
University of Michigan
Human Performance Center
330 Packard Road
Ann Arbor, MI 48109

Dr. James Howard
Dept. of Psychology
Human Performance Laboratory
Catholic University of
America
Washington, DC 20064

Dr. Lloyd Humphreys
University of Illinois
Department of Psychology
603 East Daniel Street
Champaign, IL 61820

Distribution List

Dr. Earl Hunt
Department of Psychology
University of Washington
Seattle, WA 98105

Dr. Ed Hutchins
Intelligent Systems Group
Institute for
Cognitive Science (C-015)
UCSD
La Jolla, CA 92093

Dr. Alice Isen
Department of Psychology
University of Maryland
Catonville, MD 21228

COL Dennis W. Jarvis
Commander
AFMRL
Brooks AFB, TX 78235-5601

Dr. Joseph E. Johnson
Assistant Dean for
Graduate Studies
College of Science and Mathematics
University of South Carolina
Columbia, SC 29208

CDR Tom Jones
ONR Code 125
800 M. Quincey Street
Arlington, VA 22217-5000

Mr. Daniel B. Jones
U.S. Nuclear Regulatory
Commission
Division of Human Factors Safety
Washington, DC 20555

Dr. Douglas H. Jones
Thatcher Jones Associates
P.O. Box 6680
10 Trafalgar Court
Lawrenceville, NJ 08648

Dr. Jane Jorgensen
University of Oslo
Institute of Psychology
Box 1098, Blindern
Oslo, NORWAY

Dr. Marcel Just
Carnegie-Mellon University
Department of Psychology
Schlesley Park
Pittsburgh, PA 15213

Dr. Daniel Kahneman
The University of British Columbia
Department of Psychology
#158-2053 Main Mall
Vancouver, British Columbia
CANADA V6T 1T7

Dr. Ruth Kanfer
University of Minnesota
Department of Psychology
Elliott Hall
75 E. River Road
Minneapolis, MN 55455

Dr. Demetrios Karis
Grumman Aerospace Corporation
MS C04-14
Bethpage, NY 11714

Dr. Milton S. Katz
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Steven V. Keefe
Department of Psychology
University of Oregon
Eugene, OR 97403

Dr. Wendy Kellogg
IBM T. J. Watson Research Ctr.
P.O. Box 218
Yorktown Heights, NY 10598

Dr. Scott Kelso
Haskins Laboratories,
270 Crown Street
New Haven, CT 06510

Dr. Dennis Kibler
University of California
Department of Information
and Computer Science
Irvine, CA 92717

Dr. David Kieras
University of Michigan
Technical Communication
College of Engineering
1223 E. Engineering Building
Ann Arbor, MI 48109

Dr. David Klahr
Carnegie-Mellon University
Department of Psychology
Schlesley Park
Pittsburgh, PA 15213

Dr. Ronald Knoll
Bell Laboratories
Murray Hill, NJ 07974

Dr. Sylvan Kornblum
University of Michigan
Mental Health Research Institute
205 Washtenaw Place
Ann Arbor, MI 48109

Dr. Stephen Kosslyn
Harvard University
1236 William James Hall
33 Kirkland St.
Cambridge, MA 02138

Dr. Kenneth Kotovsky
Department of Psychology
Community College of
Allegheny County
800 Allegheny Avenue
Pittsburgh, PA 15233

Dr. David M. Krantz
2 Washington Square Village
Apt. # 15J
New York, NY 10012

Dr. David B. Lambert
Naval Ocean Systems Center
Code 8411
271 Catalina Boulevard
San Diego, CA 92152-6800

Dr. Pat Langley
University of California
Department of Information
and Computer Science
Irvine, CA 92717

(Washington University/Posner) 1987/11/10

Distribution List

Dr. Marcy Lansman
University of North Carolina
The L. L. Thurstone Lab.
Davis Hall 013A
Chapel Hill, NC 27514

Dr. Jill Larkin
Carnegie-Mellon University
Department of Psychology
Pittsburgh, PA 15213

Dr. Robert Lawler
Information Sciences, FRL
GTE Laboratories, Inc.
40 Sylvan Road
Waltham, MA 02254

Dr. Paul E. Lehner
PAR Technology Corp.
7926 Jones Branch Drive
Suite 170
McLean, VA 22102

Dr. Alan M. Lesgold
Learning RAD Center
University of Pittsburgh
Pittsburgh, PA 15260

Dr. Jim Levin
Department of
Educational Psychology
210 Education Building
1310 South Sixth Street
Champaign, IL 61820-6990

Dr. John Levine
Learning RAD Center
University of Pittsburgh
Pittsburgh, PA 15260

Dr. Michael Levine
Educational Psychology
210 Education Bldg.
University of Illinois
Champaign, IL 61801

Dr. Clayton Lewis
University of Colorado
Department of Computer Science
Campus Box 430
Boulder, CO 80309

Dr. Bob Lloyd
Dept. of Geography
University of South Carolina
Columbia, SC 29208

Dr. Frederic M. Lord
Educational Testing Service
Princeton, NJ 08541

Dr. Gary Lynch
University of California
Center for the Neurobiology of
Learning and Memory
Irvine, CA 92717

Dr. Don Lyon
P. O. Box 44
Higley, AZ 85236

Dr. William L. Maloy
Chief of Naval Education
and Training
Naval Air Station
Pensacola, FL 32508

Dr. Evans Mandes
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Dr. Sandra P. Marshall
Dept. of Psychology
San Diego State University
San Diego, CA 92182

Dr. Richard E. Mayer
Department of Psychology
University of California
Santa Barbara, CA 93106

Dr. James McBride
Psychological Corporation
c/o Harcourt, Brace,
Jovanovich Inc.
1250 West 6th Street
San Diego, CA 92101

Dr. Jay McClelland
Department of Psychology
Carnegie-Mellon University
Pittsburgh, PA 15213

Distribution List

Dr. James L. McGaugh
Center for the Neurobiology
of Learning and Memory
University of California, Irvine
Irvine, CA 92717

Dr. Gail McKoon
CAS/Psychology
Northwestern University
1859 Sheridan Road
Evanston, IL 60201

Dr. Joe McLachlan
Navy Personnel RAD Center
San Diego, CA 92152-6800

Dr. James McMichael
Assistant for MPT Research,
Development, and Studies
OP 0187
Washington, DC 20370

Dr. Barbara Means
Human Resources
Research Organization
1100 South Washington
Alexandria, VA 22314

Dr. George A. Miller
Department of Psychology
Green Hall
Princeton University
Princeton, NJ 08540

Dr. Robert Mislevy
Educational Testing Service
Princeton, NJ 08541

Dr. William Montague
NPRDC Code 13
San Diego, CA 92152-6800

Mr. Melvin D. Montemarlo
NASA Headquarters
ATF-6
Washington, DC 20546

Dr. Tom Moran
Teror PARC
3333 Coyote Hill Road
Palo Alto, CA 94304

Dr. Randy Munaw
Program Manager
Training Research Division
NABRD
1100 S. Washington
Alexandria, VA 22314

Dr. Allen Munro
Behavioral Technology
Laboratories - USC
1845 S. Elena Ave., 4th Floor
Redondo Beach, CA 90277

Dr. Richard E. Mubett
University of Michigan
Institute for Social Research
Room 5261
Ann Arbor, MI 48109

Dr. Mary Jo Nissen
University of Minnesota
N218 Elliott Hall
Minneapolis, MN 55455

Deputy Technical Director
NPRDC Code 01A
San Diego, CA 92152-6800

Director, Training Laboratory,
NPRDC (Code 05)
San Diego, CA 92152-6800

Director, Manpower and Personnel
Laboratory,
NPRDC (Code 06)
San Diego, CA 92152-6800

Director, Human Factors
A Organizational Systems Lab.
NPRDC (Code 07)
San Diego, CA 92152-6800

Fleet Support Office,
NPRDC (Code 301)
San Diego, CA 92152-6800

Commanding Officer,
Naval Research Laboratory
Code 2627
Washington, DC 20390

(Washington University/Posner) 1987/11/10

Distribution List

Distribution List

Dr. Harold F. O'Neill, Jr.
School of Education - WMH 801
Department of Educational
Psychology & Technology
University of Southern California
Los Angeles, CA 90089-0031

Dr. Michael Oberlin
Naval Training Systems Center
Code 711
Orlando, FL 32813-7100

Dr. Stellan Ohlsson
Learning R & D Center
University of Pittsburgh
3939 O'Hara Street
Pittsburgh, PA 15213

Director, Research Programs,
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000

Mathematics Group,
Office of Naval Research
Code 1111MA
800 North Quincy Street
Arlington, VA 22217-5000

Office of Naval Research,
Code 1133
800 N. Quincy Street
Arlington, VA 22217-5000

Office of Naval Research,
Code 1141NP
800 N. Quincy Street
Arlington, VA 22217-5000

Office of Naval Research,
Code 1142
800 N. Quincy St.
Arlington, VA 22217-5000

Office of Naval Research,
Code 1142EP
800 N. Quincy Street
Arlington, VA 22217-5000

Office of Naval Research,
Code 1142PT
800 N. Quincy Street
Arlington, VA 22217-5000
(6 Copies)

Director, Technology Programs,
Office of Naval Research
Code 12
800 North Quincy Street
Arlington, VA 22217-5000

Special Assistant for Marine
Corps Matters,
ONR Code 00MC
800 N. Quincy St.
Arlington, VA 22217-5000

Dr. Judith Orasanu
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Jesse Orlansky
Institute for Defense Analyses
1801 W. Beauregard St.
Alexandria, VA 22311

Dr. Glenn Osge
NOSC, Code 441
San Diego, CA 92152-6800

Prof. Seymour Papert
20C-109
Massachusetts Institute
of Technology
Cambridge, MA 02139

Dr. Robert F. Pasnak
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Daira Paulson
Code 52 - Training Systems
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. James Paulson
Department of Psychology
Portland State University
P.O. Box 751
Portland, OR 97207

Dr. James W. Pellegrino
University of California,
Santa Barbara
Department of Psychology
Santa Barbara, CA 93106

Dr. Nancy Pennington
University of Chicago
Graduate School of Business
1101 E. 58th St.
Chicago, IL 60637

Dr. Ray Perez
ARI (PERI-II)
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Steven Plinker
Department of Psychology
E10-018
M.I.T.
Cambridge, MA 02139

Dr. Martha Polson
Department of Psychology
Campus Box 346
University of Colorado
Boulder, CO 80309

Dr. Peter Polson
University of Colorado
Department of Psychology
Boulder, CO 80309

Dr. Steven E. Poltrock
MCC
9430 Research Blvd.
Echelon Bldg #1
Austin, TX 78759-6509

Dr. Michael I. Posner
Department of Neurology
Washington University
Medical School
St. Louis, MO 63110

Dr. Mary C. Potter
Department of Psychology
MIT (E-10-032)
Cambridge, MA 02139

Dr. Karl Pribram
Stanford University
Department of Psychology
Bldg. 4201 -- Jordan Hall
Stanford, CA 94305

Dr. Joseph Psotka
ATM: PERI-1C
Army Research Institute
5001 Eisenhower Ave.
Alexandria, VA 22333

Dr. Mark D. Reckase
ACT
P. O. Box 168
Iowa City, IA 52243

Dr. Lynne Reder
Department of Psychology
Carnegie-Mellon University
Schenley Park
Pittsburgh, PA 15213

Dr. James A. Reggie
University of Maryland
School of Medicine
Department of Neurology
22 South Greene Street
Baltimore, MD 21201

Dr. Fred Reif
Physics Department
University of California
Berkeley, CA 94720

Dr. Daniel Reisberg
Department of Psychology
New School for Social Research
65 Fifth Avenue
New York, NY 10003

Dr. Lauren Resnick
Learning R & D Center
University of Pittsburgh
3939 O'Hara Street
Pittsburgh, PA 15213

(Washington University/Posner) 1987/11/10

Distribution List

Mr. Raymond C. Sidorosky
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Herbert A. Simon
Department of Psychology
Carnegie-Mellon University
Schenley Park
Pittsburgh, PA 15213

Dr. Zita M. Smutlis
Instructional Technology
Systems Area
ARI
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. H. Wallace Sinalco
Manpower Research
and Advisory Services
Smithsonian Institution
801 North Pitt Street
Alexandria, VA 22314

Dr. Derek Sleeman
Stanford University
School of Education
Stanford, CA 94305

Dr. Edward E. Smith
Bolt Beranek & Newman, Inc.
50 Houlton Street
Cambridge, MA 02138

Dr. Linda B. Smith
Department of Psychology
Indiana University
Bloomington, IN 47405

Dr. Robert F. Smith
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Dr. Alfred F. Smode
Senior Scientist
Code 07A
Naval Training Systems Center
Orlando, FL 32813

Distribution List

Dr. Gil Ricard
Mail Stop C04-14
Grumman Aerospace Corp.
Bethpage, NY 11714

Dr. Mary S. Riley
Program in Cognitive Science
Center for Human Information
Processing
University of California
La Jolla, CA 92093

Dr. Andrew M. Rose
American Institutes
for Research
1055 Thomas Jefferson St., NW
Washington, DC 20007

Dr. Ernst Z. Rothkopf
AT&T Bell Laboratories
Room 2D-456
600 Mountain Avenue
Murray Hill, NJ 07974

Dr. William B. Rouse
Search Technology, Inc.
25-b Technology Park/Atlanta
Norcross, GA 30092

Dr. Donald Rubin
Statistics Department
Science Center, Room 608
1 Oxford Street
Harvard University
Cambridge, MA 02138

Dr. David Rumelhart
Center for Human
Information Processing
Univ. of California
La Jolla, CA 92093

Dr. E. L. Saltzman
Haskins Laboratories
270 Crown Street
New Haven, CT 06510

Dr. Fumiko Samejima
Department of Psychology
University of Tennessee
Knoxville, TN 37916

Dr. Michael J. Smett
Perceptronics, Inc
6271 Varrel Avenue
Woodland Hills, CA 91364

Dr. Arthur Samuel
Yale University
Department of Psychology
Box 11A, Yale Station
New Haven, CT 06520

Dr. Roger Schank
Yale University
Computer Science Department
P.O. Box 2158
New Haven, CT 06520

Dr. Walter Schneider
Learning RAD Center
University of Pittsburgh
3939 O'Hara Street
Pittsburgh, PA 15260

Dr. Janet Schofield
Learning RAD Center
University of Pittsburgh
Pittsburgh, PA 15260

Dr. Hans-Wilfrid Schroll
Institut fuer Psychologie
der RWTH Aachen
Jaegerstrasse zwischen 17 u. 19
5100 Aachen
WEST GERMANY

Dr. Robert J. Seidel
US Army Research Institute
5001 Eisenhower Ave.
Alexandria, VA 22333

Dr. Michael G. Shafto
ONR Code 1142PT
800 M. Quincy Street
Arlington, VA 22217-5000

Dr. T. B. Sheridan
Dept. of Mechanical Engineering
MIT
Cambridge, MA 02139

(Washington University/Posner) 1987/11/10

Distribution List

Cdr Michael Suman, PD 303
Naval Training Systems Center
Code 451, Comptroller
Orlando, FL 32813

Dr. Steve Suoni
MIL Bldg. 31
Room 828-15
Bethesda, MD 20205

Dr. Hariharan Swaminathan
Laboratory of Psychometric and
Evaluation Research
School of Education
University of Massachusetts
Amherst, MA 01003

Mr. Brad Symson
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. John Tangney
AFOSR/ML
Boiling AFB, DC 20332

Dr. Kikumi Tatsuoka
CERL
252 Engineering Research
Laboratory
Urbana, IL 61801

Dr. Maurice Tatsuoka
220 Education Bldg
1310 S. Sixth St.
Champaign, IL 61820

Dr. Richard F. Thompson
Stanford University
Department of Psychology
Bldg. 4201 -- Jordan Hall
Stanford, CA 94305

Dr. Martin A. Tolcott
3001 Yeazey Terr., N.W.
Apt. 1617
Washington, DC 20008

Dr. Douglas Towne
Behavioral Technology Labs
1845 S. Elena Ave.
Redondo Beach, CA 90277

Dr. Robert Tsutakawa
University of Missouri
Department of Statistics
222 Math. Sciences Bldg.
Columbia, MO 65211

Dr. Michael T. Turvey
Haskins Laboratories
270 Crown Street
New Haven, CT 06510

Dr. Amos Tversky
Stanford University
Dept. of Psychology
Stanford, CA 94305

Dr. James Tweeddale
Technical Director
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Zita E. Tyler
Department of Psychology
George Mason University
4400 University Drive
Fairfax, VA 22030

Headquarters, U. S. Marine Corps
Code MPI-20
Washington, DC 20380

Dr. David Vale
Assessment Systems Corp.
2233 University Avenue
Suite 310
St. Paul, MN 55114

Dr. Kurt Van Lehn
Department of Psychology
Carnegie-Mellon University
Schenley Park
Pittsburgh, PA 15213

Dr. Jerry Vogt
Navy Personnel R&D Center
Code 51
San Diego, CA 92152-6800

Dr. Howard Walner
Division of Psychological Studies
Educational Testing Service
Princeton, NJ 08541

Distribution List

Dr. Beth Warren
Dolt Beranek & Newman, Inc.
50 Moulton Street
Cambridge, MA 02138

Dr. Norman M. Weinberger
University of California
Center for the Neurobiology
of Learning and Memory
Irvine, CA 92717

Dr. David J. Weiss
M660 Elliott Hall
University of Minnesota
75 E. River Road
Minneapolis, MN 55455

Dr. Shih-Sung Wen
Jackson State University
1325 J. R. Lynch Street
Jackson, MS 39217

Dr. Keith T. Westcott
FMC Corporation
Central Engineering Labs
1185 Coleman Ave., Box 580
Santa Clara, CA 95052

Dr. Douglas Wetzel
Code 12
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Barbara White
Dolt Beranek & Newman, Inc.
10 Moulton Street
Cambridge, MA 02238

Dr. Barry Whitsel
University of North Carolina
Department of Physiology
Medical School
Chapel Hill, NC 27514

Dr. Christopher Wickens
Department of Psychology
University of Illinois
Champaign, IL 61820

Dr. Heather Wild
Naval Air Development
Center
Code 6021
Warminster, PA 18974-5000

Dr. Robert A. Wisher
U.S. Army Institute for the
Behavioral and Social Sciences
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Martin F. Wiskoff
Navy Personnel R & D Center
San Diego, CA 92152-6800

Mr. John H. Wolfe
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. George Wong
Biostatistics Laboratory
Memorial Sloan-Kettering
Cancer Center
1275 York Avenue
New York, NY 10021

Dr. Donald Woodward
Office of Naval Research
Code 1141MP
800 North Quincy Street
Arlington, VA 22217-5000

Dr. Wallace Wulfeck, III
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Joe Yasutake
AFHRL/LRT
Lowry AFB, CO 80230

Mr. Carl York
System Development Foundation
181 Lytton Avenue
Suite 210
Palo Alto, CA 94301

Dr. Joseph L. Young
Memory & Cognitive
Processes
National Science Foundation
Washington, DC 20550

Distribution List (Washington University/Posner) 1987/11/10

AFOSR, Life Sciences Division
University of Minnesota
Department of Psychology
Minneapolis, MN 55455

Chief of Naval Education
and Training
Liaison Office
Air Force Human Resource Lab.
Operations Training Division
Williams AFB, AZ 85224

Captain P. Michael Curran
Office of Naval Research
800 N. Quincy Street
Code 125
Arlington, VA 22217-5000

Defense Technical Information Ctr.
Cameron Station, Bldg. 5
Alexandria, VA 22314
Attn: TC

Dr. Marshall J. Farr
2520 North Vernon Street
Arlington, VA 22207

Dr. John R. Frederiksen
Bolt Beranek & Newman
50 Moulton Street
Cambridge, MA 02138

Dr. Donald A. Norman
Institute for Cognitive Science
University of California
La Jolla, CA 92093

Dr. Steven Zornetzer
Office of Naval Research
Code 1140
800 N. Quincy St.
Arlington, VA 22217-5000

Dr. Jaime Carbonell
Carnegie-Mellon University
Department of Psychology
Pittsburgh, PA 15213

Dr. John J. Collins
Director, Field Research Office,
Orland
NPRDC Liaison Officer
NTSC Orlando, FL 32813

Dr. Joel Davis
Office of Naval Research
800 North Quincy Street
Code 1141NP
22217-5000

ERIC Facility Acquisitions
4833 Rugby Avenue
Bethesda, MD 20014

J. D. Fletcher
9931 Corsica Street
Vienna, VA 22180

Dr. David Navon
Institute for Cognitive Science
University of California
La Jolla, CA 92093

Dr. Robert Sasmor
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Michael J. Zyda
Naval Postgraduate School
Code 52CK
Monterey, CA 93943-5100

END

10-87

DTIC